

Buy Fioricet Online

Product Name	Price	Shipping	Total	Order
Butalbital/APAP/Caffeine (Generic Fioricet)- 50/325/40 mg- 90 Tabs	\$169	free	\$169	Order
Butalbital/APAP/Caffeine (Generic Fioricet)- 50/325/40 mg- 120 Tabs	\$199	free	\$199	Order
Butalbital/APAP/Caffeine (Generic Fioricet)- 50/325/40 mg- 180 Tabs	\$239	free	\$239	Order

Fioricet contains a combination of Acetaminophen, butalbital, and caffeine. Acetaminophen is a pain reliever and fever reducer.

Butalbital is in a group of drugs called barbiturates. It relaxes muscle contractions involved in a tension headache. Caffeine is a central nervous system stimulant. It relaxes muscle contractions in blood vessels to improve blood flow.

Fioricet is used to treat tension headaches that are caused by muscle contractions.

Fioricet may also be used for purposes not listed in this medication guide.

Butalbital is in a group of drugs called barbiturates. It relaxes muscle contractions involved in a tension headache. Caffeine is a central nervous system stimulant. It relaxes muscle contractions in blood vessels to improve blood flow.

Fioricet is used to treat tension headaches that are caused by muscle contractions.

Fioricet may also be used for purposes not listed in this medication guide.

Before taking Fioricet

Do not use Fioricet if you have used an MAO inhibitor such as furazolidone (Furoxone), isocarboxazid (Marplan), phenelzine (Nardil), rasagiline (Azilect), selegiline (Eldepryl, Emsam, Zelapar), or tranylcypromine (Parnate) in the last 14 days. A dangerous drug interaction could occur, leading to serious side effects. Tell your doctor if you have ever had alcoholic liver disease (cirrhosis) or if you drink more than 3 alcoholic beverages per day. You may not be able to take medicine that contains acetaminophen. You should not take Fioricet if you are allergic to acetaminophen, butalbital, or caffeine, or if you have porphyria.

Butalbital

Generic Name: acetaminophen/ caffeine (oral)

Brand Names: Esgic, Esgic-Plus, Fioricet

HOW TO USE: To prevent upset stomach, take this medication with food or milk. Pain medications work best in preventing pain before it occurs. Once the pain becomes intense, the medication is not as effective in relieving it. Take this as prescribed. Do not increase your dose or take it more frequently without your doctor's approval.

SIDE EFFECTS: May cause lightheadedness, dizziness, drowsiness, stomach upset, nausea, and sleeping disturbances the first few days as your body adjusts to the medication. If these symptoms persist or become bothersome, inform your doctor. Notify your doctor immediately if you develop: a rapid/ irregular heartbeat. An allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS: Tell your doctor your medical history, especially of: serious liver disease, alcohol or narcotic dependence, emotional/mental conditions, heart disease (arrhythmias, recent MI), stomach/intestinal ulcers, any allergies. Acetaminophen may cause liver damage. Daily use of alcohol, especially when combined with acetaminophen, may increase your risk for liver damage. Check with your doctor or pharmacist for more information. To prevent oversedation, avoid using alcohol and other sedative type medications while taking this. This drug may make you dizzy or drowsy; use caution engaging in activities requiring alertness such as driving or using machinery. Avoid alcoholic beverages. This medication contains caffeine. Avoid drinking large amounts of caffeinated beverages while taking this medication as excessive nervousness and irritability can occur. This medication should be used during pregnancy only if clearly needed. Small amounts of this medication appear in breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS: Tell your doctor of all prescription and nonprescription drugs you may use, especially of: "blood thinners" Tell your doctor or pharmacist of all

prescription and nonprescription drugs you may use that cause drowsiness such as: medicine for sleep (e.g., sedatives), tranquilizers, anti-anxiety drugs (e.g., diazepam), narcotic pain relievers (e.g., codeine), psychiatric medicines (e.g., phenothiazines such as chlorpromazine, or tricyclics such as amitriptyline), anti-seizure drugs (e.g., carbamazepine), muscle relaxants, antihistamines that cause drowsiness (e.g., diphenhydramine). Because this medication contains acetaminophen, check with your doctor or pharmacist before taking other medications containing acetaminophen which is commonly found in nonprescription pain relievers and cough-and-cold products. Read labels carefully to check ingredients. This drug may interfere with the effectiveness of birth control pills. Discuss using other methods of birth control with your doctor. Do not start or stop any medicine without doctor or pharmacist approval.


OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include vomiting, unusual drowsiness, lack of feeling alert, slow or shallow breathing, cold or clammy skin, loss of consciousness, dark urine, stomach pain, and extreme fatigue.

NOTES: Do not share this product with others.

MISSED DOSE: If you miss a dose, take it as soon as remembered; do not take if it is almost time for the next dose, instead, skip the missed dose and resume your usual dosing schedule. Do not “double-up” the dose to catch up.

STORAGE: Store at room temperature away from moisture and sunlight. Do not store in the bathroom

Buy Fioricet Online in US Pharmacy



A lot of American suffer from all kinds of headaches. Fioricet contains a combination of acetaminophen, butalbital, and caffeine. Acetaminophen is a pain reliever and fever reducer. Butalbital is in a group of drugs called barbiturates. It relaxes muscle contractions involved in a tension headache. Caffeine is a central nervous system stimulant. It relaxes muscle contractions in blood vessels to improve blood flow.

Fioricet is prescribed for tension headache. But a lot of patients take fioricet for migraine headache and cluster headache. Actually more patients take fioricet for migraine headache than for tension headache. More than 90% patients think fioricet is good and effective for tension headache and more than 80% patients think fioricet is good and effective for Migraine Headache.

Some patients even use fioricet together with gabapentin for cancer pain relieving. AMORITZ, 45-54 Female, reviewed fioricet "I have been diagnosed with cervical cancer and the pain is severe where I'm squirming on the floor needing to go to the ER, but Fioricet along with Gabapentin work hand in hand and I am 100% pain free. This medicine goes directly to my nerve endings and all pain is gone. I wish the Cancer would be gone as well."

AMORITZ is right, Fioricet contains a combination of acetaminophen, butalbital, and caffeine. Acetaminophen is a pain reliever and fever reducer. Butalbital is in a group of drugs called barbiturates. It relaxes muscle contractions involved in a tension headache. Caffeine is a central nervous system stimulant. It relaxes muscle contractions in blood vessels to improve blood flow. So Fioricet is good to relieve nerve pain together with Gabapentin. For more Fioricet and Fioricet Side Effects information, please check [Fioricet](#).



Fioricet generic is not controlled but the best seller tramadol is controlled by DEA. Some patients are wondering why Fioricet is not controlled but Fiorinal is controlled. The reason is that Fiorinal contains aspirin but Fioricet only contains non-aspirin pain killer and barbiturate butalbital. The potential for abuse is decreased if a sufficient amount of a non-controlled substance (e.g., aspirin or acetaminophen) is added. The ratio required for exemption is 70 mg of acetaminophen to 15 mg of butalbital. In contrast, the ratio of aspirin to butalbital is 188 mg to 15 mg. The acetaminophen to butalbital ratio is approximately 97 mg to 15 mg in Fioricet, thus it is exempt from being controlled. Another reason is that drug abusers can easily perform an "acid wash" to remove the aspirin (acetylsalicylic acid) component in Fiorinal but it is much more difficult to remove the acetaminophen component in Fioricet.

For more Fioricet information, the best place is to read the articles in rxlists.com and [Fioricet](#).

Usual Adult Dose of Fioricet for Headache:

Acetaminophen 325 mg, butalbital 50 mg, and caffeine 40 mg: 1 or 2 tablet(s), capsule(s), or tablespoonful(s) orally every 4 hours. Maximum daily dose: 6 doses. Please never exceed your max dosage. If you overdose your Fioricet pills, you will hurt your liver, kidney, and heart. Hurting is caused by acetaminophen (Tylenol). The safe acetaminophen dosage is 3000mg per day. Healthy adults taking regular doses of up to 4,000 mg a day show little evidence of toxicity (although some researchers

disagree). In 2011 the U.S. Food and Drug Administration launched a public education program to help consumers avoid overdose, warning: "[Acetaminophen can cause serious liver damage if more than directed is used. Some doctors even suggest acetaminophen be controlled or become prescription.](#)"

Butalbital is one kind of barbiturate. The average daily dose for the barbiturate addict is usually about 1500 mg. You cannot be addictive on butalbital (totally 240mg per day taken) in Fioricet. But you may depend on butalbital in Fioricet. Major withdrawal symptoms (convulsions and delirium) may occur within 16 hours and last up to 5 days after abrupt cessation of these drugs. Intensity of withdrawal symptoms gradually declines over a period of approximately 15 days.

Usual Pediatric Dose of Fioricet for Headache:

12 years and older:

Acetaminophen 300 mg, butalbital 50 mg, and caffeine 40 mg:

1 or 2 capsule(s) orally every 4 hours as needed. Maximum daily dose: 6 doses.

Who cannot Buy Fioricet Online ?

Some patients cannot take Fioricet. If you have liver, kidney, or heart diseases, please do not take Fioricet. Because Fioricet contains barbiturate butalbital, it can cross the placental barrier. We do not suggest you order Fioricet online if you are pregnant or in breast feeding period.

If you are pregnant, please go to your street doctors and let them to check whether you are OK to take Fioricet. If you are addictive to alcohol or other pain killer medicines, please do not buy Fioricet online, you must find a street doctor to solve your addictive first before you order Fioricet, or buy Fioricet online. You must complete the health questionnaires very carefully because the US licensed doctors and pharmacists will review your health questionnaires to approve your prescription or not.

For more detail Fioricet prescription instruction, you can check [Wiki Fioricet](#). The pharmacy will also send you a instruction on Fioricet usage and introduction. Please be very careful to take the Fioricet and tramadol within the max dosage. Never exceed the max dosages. Or you will be seriously damaged.

